

AIPPG Paper 2002 (Part 13 of 25)

1. Which of the following is true
 - a. Imipramine is used in the treatment of Endogenous depression
 - b. Diazepam has lesser sedative propensity as compared to barbiturates
 - c. Fluvoxamine is associated with weight gain
 - d. Thioridazine causes lesser extrapyramidal side effects

2. Drugs used in the treatment of Schizophrenia include
 - a. Chlorpromazine
 - b. Haloperidol
 - c. Olanzapine
 - d. Imipramine
 - e. Risperidone

3. Schizophrenia has which of the following features in common with depression
 - a. Formal thought disorder
 - b. Social withdrawal
 - c. Poor personal care
 - d. Decreased interest in sex
 - e. Inappropriate behavior

4. Suicidal tendencies are seen in
 - a. Schizophrenia
 - b. Post traumatic stress disorder
 - c. Depression
 - d. OCD
 - e. Anxiety disorder

5. Features seen in Obsessions and compulsions is/are
 - a. Repetetiveness
 - b. Irresistiblility
 - c. Unpleasant
 - d. Social withdrawal
 - e. Poor personal care

6. Pleural fibrosis is caused by
 - a. Phenytoin
 - b. Methysegide
 - c. Amiodarone
 - d. Ergotamine
 - e. Ranitidine

7. True regarding bromocriptine
 - a. Natural derivative
 - b. Synthetic derivative
 - c. Also has alpha blocking property
 - d. Decreases GI motility
 - e. Acts on both D1 and D2

8. A lady with 16 weeks pregnancy presents with acute appendicitis. Management includes
 - a. Conservative treatment
 - b. Do early surgery
 - c. Appendectomy with termination of pregnancy
 - d. Appendectomy following child birth after 3rd trimester
 - e. Medical treatment with surgery on recurrence

9. A lady with 4 months ammenorhea presents with pain abdomen, constipation, and vomiting. What will be done in her management
 - a. X ray abdomen

Visit examrace.com for free study material, doorsteptutor.com for questions with detailed explanations, and "Examrace" YouTube channel for free videos lectures

- b. USG
- c. CT scan
- d. Complete obstetrical evaluation

10. True regarding hirshsprungs disease is/are

- a. Seen in infants and children only
- b. Absence of auerbachs plexus in the involved segment
- c. The involved segment is the dilated colon
- d. Bleeding per rectum is a presenting feature
- e. Surgery is used in the treatment