

MAHE PG Solved Paper 2003 (Part 14 of 19)

1. The lymphatic spread is seen in all of the following except

- a. Terratoma
- b. Seminoma
- c. Squamous cell carcinoma
- d. Basal cell carcinoma

Answer: d

2. Which of the following is true regarding ESR

- a. Increased in polycythemia
- b. Increased in sickle cell anemia
- c. Useful in monitoring the treatment response in temporal arteritis
- d. Decreased in pregnancy

Answer: c

3. Which of the following is a T cell malignancy

- a. CML
- b. Hodgkins disease
- c. Mycosis fungoides
- d. Burkits lymphoma

Answer: c

4. Autoimmune haemolytic anemia is seen in which of the following conditions commonly

- a. Hairy cell leukaemia
- b. Chronic myeloid leukaemia
- c. Chronic lymphoid leukaemia
- d. Lymphomas

Answer: b

5. The philadelphia chromosome is classically associated with which of the following

- a. CLL
- b. CML
- c. Burkits lymphoma
- d. Hairy cell leukaemia "AI PP G" c om

Answer: b

6. All of the following are poor prognosticators except

- a. Mediastinal tumors
- b. Leucocyte count > 100000 cells/cmm at the time of presentation
- c. Age between 2 – 10 years
- d. Aneuploidy

Answer: c

7. The treatment of choice in pancreatic abcess is

- a. Cystogastrostomy
- b. Cystoenterostomy
- c. External drainage
- d. Antibiotics

Answer: c

8. Chronic pancreatitis is associated with all of the following except

- a. Calcification of pancreatic duct
- b. Steatorrhea
- c. Bronze skin discoloration
- d. Diabetes mellitus

Answer: c

9. Pyloric stenosis is associated with all of the following except

- a. Weight loss

Visit examrace.com for free study material, doorsteptutor.com for questions with detailed explanations, and "Examrace" YouTube channel for free videos lectures

- b. Diarrhea
- c. Ramsteads operation
- d. Hypokalemia is seen

Answer: b

10. Functions of the amniotic fluid include

- a. Decreased fetal weight
- b. Reduce trauma to fetus
- c. Conjugate and store hormones
- d. All of the above

Answer: b