

## NIMHANS PG Paper 2002 (Part 5 of 14)

1. Which of the following is true
  - a. The right gastric artery arises from the common hepatic artery
  - b. Ligation of the splenic artery causes infarction of the greater curvature of the stomach
  - c. Gastroepiploic arteries arise from gastroduodenal and splenic arteries
  - d. The lesser curvature gets its blood supply mainly from the short gastric vessels
  
2. A smoker presented with altered sensorium. His blood osmolality was found to be 240 and urine osmolality 340. His Na<sup>+</sup> is 122 meq/L. The diagnosis is
  - a. Diabetes insipidus
  - b. SIADH
  - c. Renal tubular acidosis
  - d. Cerebellar degeneration
  
3. A patient came with history of fall and on examination there was tenderness between the extensor pollicis longus and brevis. The likely lesion is
  - a. Scaphoid
  - b. 1st metacarpal
  - c. Lower end of radius
  - d. Trapezoid
  
4. A child when playing had cat scratches over his hand. After 2 days he developed lymphadenopathy. Which group of the lymphnodes are involved if the scratches are over the region where the cephalic vein penetrates into the deeper fascia.
  - a. Deltopectoral group
  - b. Lateral group
  - c. Clavipectoral group
  - d. Central group

5. Inability to abduct left eye with LMN fascial palsy on same side. The lesion is in
- Left pons
  - Right pons
  - CP angle tumor
  - Cerebellar lesions
6. In a patient with burns
- Carbon monoxide is highly toxic to the pulmnory tract
  - Early tracheostomy is indicated because it is difficult to to pass tube through inflamed pharynx
  - The laryngeal edema can occur as late as 48 hours
7. In a patient presenting odynophagia with oral thrush the chest xray was found to be normal. The CD4 count was found to be 2400. The next step in management is
- Fluconazole
  - Acyclovir
  - Endoscopy with biopsy of lesion
  - IV amphotericin B
8. Pulmonary fibrosis is caused by all of the following except
- Clindamycin
  - Nitrofurantoin
  - Methysergide
  - Amidarone
9. A drug which causes parkinsonism and is not to be used in patients with this type of disease is
- Methysergide
  - Haloperidol
  - Riluzole
  - Phenytoin

10. In a child with hernia on the right side the management is

- a. High ligation of the lesion is done
- b. The distal sac needs to be excised as it can develop hydrocele
- c. Opposite side needs to be explored as bilaterality is common
- d. X ray of the abdomen is useful for diagnosis