

Physics MCQs for Competitive Exams Part 8

Question:

If young's double slit experiment is performed in water.

1. The fringe width will decrease
2. The fringe width will increase
3. The fringe width will remain unchanged
4. There will be no fringe

Question:

The first diffraction minimum due to single slit diffraction is θ , for a light of wave length 5000\AA . If the width of the slit is $1 \times 10^{-4} \text{ cm}$. then the value of θ is

1. 300
2. 450
3. 600
4. 150

Question:

Non-coherent sources emit light beam of intensities I and $4I$. the maximum and minimum intensities in the resulting beam are

1. $9I$ and $3I$
2. $9I$ and $5I$
3. $5I$ and I
4. $5I$ and $3I$

Question:

Light propagates 2 cm distance in glass of refractive index 1.5 in time t_0 . In the same time t_0 , light propagates a distance of 2.25 cm in a medium. The refractive index of the medium is

Visit examrace.com for free study material, doorsteptutor.com for questions with detailed explanations, and "Examrace" YouTube channel for free videos lectures

1. $\frac{4}{3}$
2. $\frac{3}{2}$
3. $\frac{8}{3}$
4. *None of these*

Question:

Two wave fronts are emitted from coherent sources of path difference between them is 2.1 micron. Phase difference between the wave fronts at that point is 7.692π Wave length of light emitted by source will be

1. 5386 Å
2. 5400 Å
3. 5460 Å
4. 5892 Å

Question:

A spherical air bubble in water will act as

1. Convex lens
2. Concave lens
3. Glass plate
4. Plano convex lens

Question:

A concave lens can be used as a simple magnifier if the object lies

1. Beyond f
2. Within the focal length
3. Between f and 2f
4. At 2f

Question:

Visit examrace.com for free study material, doorsteptutor.com for questions with detailed explanations, and "Examrace" YouTube channel for free videos lectures

For an equilateral prism the angle of minimum deviation is 30° . Then the refractive index of the material of the prism is

1. $\frac{1}{2}$
2. 2
3. 4
4. $\sqrt{2}$