

Q.1 — 30 Carry One Mark Each.

- 1. The structural feature common for propranolol, atenolol, pindolol, metoprolol in the side chain is**
(A) Isopropylamino propan-2-ol
(B) Dimethylamino propan-2-ol
(C) Diethylamino propan-2-ol
(D) Dibutylamino propan-2-ol
- 2. When N-methyl group of morphine is replaced with an allyl group, the compound formed is**
(A) Naloxone-morphine antagonist
(B) Naltrexone-morphine agonist
(C) Nalorphine-morphine antagonist
(D) Nalbuphine-morphine agonist/antagonist
- 3. Nitrazepam can be synthesized from**
(A) 2-Bromo-5-amino benzophenone
(B) 2-Nitro-2-chloro acetophenone
(C) 2-Amino-5-nitro cyclohexanone
(D) 2-Amino-5-nitro benzophenone
- 4. Clavulanic acid has a beta lactam ring fused to**
(A) Thieryl system
(B) Thiadiazole system
(C) Thiazolidine system
(D) Oxazolidine system
- 5. A drug which has antipyretic, anti-inflammatory and antiplatelet activity is**
(A) Sulfinpyrazone
(B) Aspirin
(C) Ticlopidine
(D) Acetaminophen
- 6. Wild cherry bark contains prunasin which is a**
(A) Phenolic glycoside
(B) Isothiocyanate glycoside
(C) Coumarin glycoside
(D) Cyanoglucoside glycoside
- 7. Euphonia, Euphorbia and Ephedra equisetina can be distinguished by type of**
(A) Breeding
(B) Stamens
(C) Fleshy leaves
(D) Alkaloids
- 8. Micropropagation of the plants is carried out through**
(A) Cross fertilization
(B) Seed germination
(C) Plant tissue culture

(D) Grafting

9. Aconitine belongs to the group of

- (A) Steroidal alkaloids
- (B) Terpenoidal alkaloids
- (C) Indole alkaloids
- (D) Quinoline alkaloids

10. Crude fiber value of a drug is a measure of

- (A) Soft tissue matter
- (B) Woody matter
- (C) Mineral matter
- (D) Organic matter

11. One of the units used for expressing pressure is 'torr' and it is equivalent to

- (A) cm of Hg
- (B) mm of Hg
- (C) psi
- (D) gauss

12. Removal of a single electron from a molecule results in the formation of

- (A) Fragment ion
- (B) Metastable ion
- (C) Molecular ion
- (D) Rearrangement ion

13. Nuclear magnetic moment is NOT known by

- (A) ^{13}C
- (B) ^{16}O
- (C) ^1H
- (D) ^{15}N

14. Derivatisation techniques in HPLC are intended to enhance

- (A) Molecular weight
- (B) Detectability
- (C) Reversibility
- (D) Reproducibility

15. A conductance cell is calibrated by using a solution of known conductivity, i.e., usually a solution of

- (A) NaCl
- (B) Hg_2Cl_2
- (C) KCl
- (D) Na_2SO_4

16. Metoclopramide is generally used for

- (A) Prophylaxis of vomiting
- (B) Preventing motion sickness
- (C) Treating irritable bowel syndrome

(D) Treatment of pancreatic insufficiency

17. DNA amplification by the polymerase chain reaction uses

- (A) Thermus aquaticus DNA polymerase
- (B) DNA topoisomerase
- (C) RNA polymerase
- (D) DNA helicase

18. Identify the non-pathogenic organism

- (A) *Mycobacterium bovis*
- (B) *Mycobacterium smegmatis*
- (C) *Mycobacterium avium*
- (D) *Mycobacterium intracellulare*

19. Bioassays are carried out to

- (A) Measure the pharmacological activity of a drug
- (B) Avoid clinical trials for new drugs
- (C) Detect the impurity in a given drug
- (D) Screen for pharmacogenetic influences of new drugs.

20. A direct way of studying idiosyncratic reactions to a given drug is by

- (A) changing the route of drug administration
- (B) changing the assay method
- (C) pharmacogenomics
- (D) structure activity relationship studies of a family of compounds

21. An example of haemopoietic growth factor is

- (A) platelet derived growth factor
- (B) epidermal growth factor
- (C) iron dextran
- (D) erythropoietin

22. Saffranin is used as a reagent to detect

- (A) Gram-negative bacteria
- (B) Gram-positive bacteria
- (C) Acid fast bacteria
- (D) Myxozoa

23. Sulphonamides do not have adverse drug interaction with

- (A) Oral anticoagulants
- (B) Sulfamylon (a hypoglycemic agents)
- (C) Hydralazine (anticonvulsants)
- (D) Dihydrofolate reductase inhibitors

24. Simvastatin belongs to

- (A) HMG CoA reductase inhibitor type of antilipidemic agents
- (B) HMG CoA reductase inhibitor type of anticoagulant agents
- (C) Fibrate type of anticoagulant agents

(D) Fibrate type of antilipidemic agents

25. HIV infection can be clinically controlled with

- (A) Cytarabine
- (B) Acyclovir
- (C) Zidovudine
- (D) Amantadine

26. The measure of cohesive strength of the cross linking that occurs between gelatin molecules and is proportional to the molecular weight of gelatin is called

- (A) Bloom strength
- (B) Viscosity
- (C) Surface tension
- (D) Partition coefficient

27. A water soluble substance used as coating material in microencapsulation process is:

- (A) Polyethylene
- (B) Silicone
- (C) Hydroxy ethyl cellulose
- (D) Paraffin

28. One of the following is used as a solubilizing agent to solubilize testosterone in pharmaceutical liquid dosage forms.

- (A) Sucrose monoesters
- (B) Lanolin esters
- (C) Lanolin ethers
- (D) Tweens

29. One of the following is used as a pH dependent controlled release excipient.

- (A) Carnauba wax
- (B) Hydroxy propyl methyl cellulose phthalate
- (C) Methyl cellulose
- (D) Glyceryl monostearate

30. The Schedule in D & C act that deals with the standards for disinfectant fluids is:

- (A) Schedule B
- (B) Schedule F
- (C) Schedule G
- (D) Schedule M

Q.1 – 20 Carry Two Marks Each.

31. The carboxyl group of aspirin is esterified with N-acetyl-p-aminophenol to get

- (A) 4-acetamidophenyl-O-acetyl salicylate
- (B) 4-Acetamidophenyl-O-acetyl salicylate
- (C) O-(2-hydroxy benzoyl) salicylic acid
- (D) 2-acetamidophenyl-O-acetyl salicylate

32. JUPAC system of nomenclature for diclophenac sodium (BP) is

- (A) Sodium 2-[(2, 6-Dichlorophenyl) amino] phenyl acetate
- (B) Sodium 3-[(2, 6-Dichlorophenyl) amino] phenyl acetate
- (C) Sodium 2-[(2-Chlorophenyl) amino] phenyl acetate
- (D) Sodium 2-[(6-Chlorophenyl) amino] phenyl acetate

33. 1-(2-Aminoethyl) perhydrozocine on treatment with S-methyl isothiurea gives rise to an adrenergic neuron blocking agent

- (A) Bethanidine
- (B) Mecamylamine
- (C) Guanadrel
- (D) Gauenthidine

34. Quercetin is

- (A) 5, 7, 3-Trihydroxy flavone
- (B) 5, 7, 3, 4-Tetrahydroxy flavone
- (C) 3, 5, 7, 3, 4-Pentahydroxy flavonol
- (D) 3, 5, 7, 3, 4-Pentahydroxy flavanone

35. Meconic acid is a chemical marker for the genus

- (A) Piper
- (B) Pilocarpus
- (C) Prunus
- (D) Papaver

36. A novel diterpenoid isolated from the bark of *Taxus brevifolia* is

- (A) Demecolcine
- (B) Pacitaxel
- (C) Vinblastin
- (D) Brevifolcin

37. The absorption maximum for polar compounds is usually shifted with change in polarity of the solvents due to

- (A) Hydrogen bonding
- (B) Chemical reaction
- (C) Ionization of the compounds
- (D) Change in dielectric permittivity

38. A titration in which potential applied across two electrodes is maintained at a constant value and the current is measured and plotted against volume of titrant is

- (A) Potentiometric titration (B) Amperometric titration
- (C) Displacement titration (D) Conductometric titration

39. The parameter in the elution curve that is proportional to the concentration of a compound in gas chromatographic effluent is the

- (A) Number of peaks
- (B) Width of the peak
- (C) Area under the peak
- (D) Shape of the peak

40. A drug solution has a half life of 21 days. Assuming that the drug undergoes first order kinetics, how long will it take for the potency to drop to 90% of the initial potency?

- (A) 3.2 days
- (B) 9.6 days
- (C) 16 days
- (D) 6.4 days

41. An amphiphilic surfactant used in pharmaceutical dispense system is:

- (A) Bile salts
- (B) Lecithin
- (C) Sorbitan monolaurate
- (D) Sorbitan monostearate

42. An abrasive used in dentifrices is

- (A) Dicalcium phosphate
- (B) Sodium carboxy methyl cellulose
- (C) Sodium lauryl sulfate
- (D) Dioctyl sodium sulfosuccinate

43. An electrochemical method that enhances the transport of some solute molecules by creating a potential gradient through the skin tissue with the applied electrical current or voltage is called

- (A) Electrophoresis
- (B) Iontophoresis
- (C) Osmosis
- (D) Implants

44. A patient with rheumatoid arthritis has been taking acetyl salicylic acid regularly. However, recently she has been experiencing skin rashes, swelling and pain due to salicylate resistance. She has occult blood in her feces. Suggest an appropriate drug suitable for her from those mentioned below:

- (A) Paracetamol
- (B) Celecoxib
- (C) Piroxicam
- (D) Naproxen

45. The break down of fibrin is catalyzed by

- (A) Plasmin
- (B) Rennin
- (C) Urokinase
- (D) Streptokin

46. Which one of the these best describes a process carried out to render a drug pharmaceutically more acceptable?

- (A) Enteric coating of diclofenac.
- (B) Co-administration of aspirin with antacids.
- (C) Use of colloidal suspensions or liposomes for administering amphotericin-B.
- (D) Synthesis of an analogue to obtain high receptor specificity.

47. Azithromycin is clinically administered once daily as compared to erythromycin which is administered every 6 hours because, azithromycin

- (A) Penetrates into most tissues and is released very slowly.
- (B) Has a methylated nitrogen in its lactone ring which renders it much more potent than erythromycin.
- (C) Is a very potent antibiotic but not tolerated well in the gastrointestinal tract.
- (D) Is usually presented in a sustained release dosage form.

48. A patient showing muscle rigidity, bradykinesia, tremors and postural instability is administered levo-dopa. Which of the properties of levo-dopa is not true?

- (A) Levo-dopa is preferred over dopamine because it can cross the blood brain barrier.
- (B) Levo-dopa is the levorotatory stereoisomer of 3, 4-dihydroxy phenethylamine.
- (C) Levo-dopa gets decarboxylated in the brain to dopamine.
- (D) Levo-dopa is administered because of its strong antagonistic action on dopamine receptors.

49. Autoimmunity refers to

- (A) an automatic trigger of the immune system directed against a specific pathogen.
- (B) failure to distinguish between self and non-self.
- (C) an automatic segregation of T and B cells.
- (D) failure of B-cells to interact with T-cells.

50. Which of these is true about the discovery of HB antigen in the blood of people infected with Hepatitis-B?

- (A) It provided a basis for vaccine design.
- (B) It indicated that specific vaccine containing HB antigen can be designed for Hepatitis-B.
- (C) It has not been of much significance.
- (D) It indicated that Hepatitis is not a viral disease.

51. Which drug molecule does NOT have phenylethyl amine moiety?

- (A) Amphetamine
- (B) Glyburide
- (C) Pheniramine
- (D) Mescaline

Q.52-58 are multiple selection items. P, Q, R, S are the options. Two of these options are correct. Choose the correct combination among A, B, C and D.

52. There are two methods by which the duration of action of insulin may be prolonged.

- (1) Binding with resins.
 - (2) Esterification of amino acid residues.
 - (3) Forming of complex of insulin with protein.
 - (5) Modification of particle size.
- (A) Q, R (B) R, S (C) P, S (D) P, R

53. The attributes of cycloserina are

- (P) No tautomerism shown.
 - (Q) Exists in equilibrium with its tautomeric enolic form.
 - (R) Stable in alkaline solution, destroyed rapidly at neutral or acidic pH.
 - (S) Stable in neutral solution, destroyed in alkaline pH.
- (A) R, S (B) P, Q (C) Q, R (D) P, R

54. Compared to benzyl penicillin, amoxicillin has the following advantages in biological properties.

- (P) The amino group renders the antibiotic resistant to acid catalysed degradation.
 - (Q) The spectrum of acidity is broadened.
 - (R) The amino group of renders penicillinase resistance to the compound.
 - (S) The phenolic group renders penicillinase resistance to the compound.
- (A) P, Q (B) P, R (C) P, S (D) Q, R

55. The identification of propellants in pharmaceutical aerosols is carried out by

- (P) Gas-chromatography
 - (Q) Tag-open cup apparatus
 - (R) Pyknometer
 - (S) JR Spectrophotometer
- (A) P, Q (B) P, S (C) Q, R (D) R, S

56. Schedule 'H' and Schedule 'S' as per the Drugs and Cosmetics Act deal with the following.

- (P) Prescription drugs which are required to be sold only retail only on prescription of R.M.P.
 - (Q) Standards for cosmetics
 - (R) Biological and special products
 - (S) List of coal tar colours permitted to be used in cosmetics and soaps
- (A) P, Q (B) P, R (C) Q, S (D) R, S

57. Myristica fragrans Houtt. has two of the following characteristics.

- (P) An indeciduous tree, which produces drupaceous, pale yellow fruits.
 - (Q) Each fruit has several round seeds with smooth surface and lignaceous tegument, and the orange red fleshy aril — the mace, is present inside the seed.
 - (R) A deciduous tall tree, which produces lignaceous capsules.
 - (S) Each fruit has a unique aroid seed, with lignified tegument, surrounded by orange red lacinate fleshy aril — the mace.
- (A) Q, R (B) P, R (C) S (D) Q, S

58. In size exclusion chromatography the stationary phases used are:

- (P) Alumina (D)
 - (Q) Dextran (R)
 - (S) Agarose (S)
 - (T) Styrene
- (A) P, S (B) Q, R (C) Q, S (D) P, R

59-65 ARE 'MATCHING' exercises. Match Group I with Group II. Choose the correct combination among the alternatives A, B, C and D.

Group II

Intermediates from which Group I drugs are synthesized

(1) Aziridin and thiophosphoryl chloride

(Q)

(2) 4-Chlorophenol

(R)

(3) 4-Chlorobenzhydryl chloride

(S) Alprazolam

(4) 2-Amino-5-Chloro benzophenone

Codes:

(A) P - 3 Q - 2 R - 1 S - 4

(B) P - 4 Q - 2 R - 1 S - 3

(C) P - 2 Q - 4 R - 3 S - 1

(D) P - 1 Q - 2 R - 4 S - 3

60

Group I

Cardiac Agents

(P) Digitoxin

(Q)

Dobutamine

(R) Sotalol

(S) Nicardipine

Group II

Mechanism of action

(1) Produces negative inotropic effect by blocking calcium channels

(2) Depresses adrenergically enhanced calcium influx through beta receptor blockade

(3) Causes elevation of cAMP levels by stimulation of adenylate cyclase

(4) Inhibits membrane bound sodium potassium ATPase pump.

Codes:

(A) P-4Q-3R-2S-1

(B) P-3Q-4R-1S-2

(C) P-4Q-2R-3S-1

(D) P-4Q-3R-1S-2

61.

Group I

Technique employed

(P) Visible spectrophotometry

(Q) IR spectrophotometry

(R) NMR spectrophotometry

(S) Fluorescence

Group II

Source of Radiation

(1) RF Source transmitter

(2) Xenon lamp

(3) Tungsten lamp

(4) Nernst glower

spectrophotometry

Codes:

- (A) P - 2 Q - 4 R - 3 S - 1
- (B) P - 3 Q - 2 R - 1 S - 4
- (C) P - 3 Q - 4 R - 1 S - 2
- (D) P - 4 Q - 1 R - 3 S - 2

62.

Group I

Amino acids

Common degradative products that are citric acid cycle intermediates or their precursors.

- (P) Aspartic acid (1) Succinyl CoA
- (Q) Arginine (2) Alpha-Ketoglutarate
- (R) Serine (3) Fumarate
- (S) Methionine (4) Pyruvate

- (A) P - 3 Q - 2 R - 4 S - 1
- (B) P - 3 Q - 1 R - 4 S - 2
- (C) P - 1 Q - 2 R - 3 S - 4
- (D) P - 4 Q - 2 R - 3 S - 1

63.

Group J
Tablet
defects

Group JJ Explanations

- (P) Picking (1) A term used to describe the surface material from a tablet that is sticking to and being removed from the tablet's surface by a punch.
- (Q) Sticking (2) Term refers to tablet material adhering to the die wall.
- (R) Mottling (3) Term refers to an unequal distribution of colour on a tablet.
- (S) Lamination (4) Term refers to separation of a tablet into two or more distinct layers.

Codes:

- (A) P - 3 Q - 4 R - 1 S - 2
- (B) P - 4 Q - 1 R - 2 S - 3

64.

- | | |
|------------------|-------------------|
| Group I | Group II |
| Lanatosides | Aglycone |
| (P) Lanatoside A | (1) Gitaxigenin |
| (Q) Lanatoside B | (2) Diginatigenin |
| (R) Lanatoside C | (3) Digoxigenin |
| (S) Lanatoside D | (4) Digitoxigenin |

- (A) P - 1 Q - 2 R - 3 S - 4
(B) P - 3 Q - 1 R - 2 S - 4
(C) P - 2 Q - 4 R - 3 S - 1
(D) P - 3 Q - 3 R - 1 S - 4

65.

- | | |
|--------------------------------|-------------------|
| Group I | Group II |
| Specific chemical test | Phytoconstituents |
| (P) Thalleioquin Test | (1) Hyoscyamine |
| (Q) Murexide Test | (2) Barbaloin |
| (R) Vitali-Morin Test | (3) Quinine |
| (S) Modified Borntrager's Test | (4) Theophylline |

- (A) P - 4 Q - 1 R - 3 S - 2
(C) P - 3 Q - 4 R - 2 S - 1
(B) P - 1 Q - 3 R - 4 S - 1
(D) P - 2 Q - 2 R - 3 S - 4

Data for Q.66 are based on the statements/problem. Choose the correct answer for each question from among the options A, B, C and D.

Data for question 66 to 68:

In a formulation development laboratory a tablet is to be formulated. The core tablet has a bad taste and requires physical and chemical protection of the drug from moisture. The tablet should also deliver the drug for local action in the intestine.

Which suggests a suitable method.

- (A) Sugar coating
(B) Film coating
(C) Enteric coating
(D) Sub coating

67. Choose the correct coating material to be used.

- (A) Sugar
- (B) Acacia
- (C) Ethyl cellulose
- (D) Cellulose acetate phthalate

68. Choose the correct solvent for the coating material.

- (A) Acetone
- (B) Water
- (C) Propylene glycol
- (D) Glycerine

Data for questions 69 and 70:

Compound A with the formula C_2H_7N shows the following important bands in the IR spectra; (a) 3423 cm^{-1} (b) 3236 cm^{-1}

69. Assign these bands to the important group in the compound A.

- (A) $-CH_3$ (B) $-NH$ (C) $-CN$ (D) $=C=N$

70. On treatment with nitrous acid the compound A is converted to B, which shows a strong band at 3430 cm^{-1} . Assign the absorption band for the group formed in the product.

- (A) $-OH$ (B) $=C=N$ (C) $-COOH$ (D) $-N=N$

Data for questions 71 to 73:

In the assay of sulfamethoxazole ($C_{10}H_{11}N_3O_2S$), 0.2 g of the sample was dissolved in 50 ml of 2M HCl. To this was added 3 g of NaOH and the titration was carried out.

71. Titration was carried out using

- (A) $NaNO_2$ to estimate the amino group.
- (B) $NaNO_2$ to estimate the sulphohamido group.
- (C) $NaOH$ to estimate the amino group.
- (D) $NaOH$ to estimate the sulphohamido group.

72. The end point in the assay was determined by

- (A) Conductometric method
- (B) Using an indicator
- (C) Potentiometric method
- (D) Photometric method

73. If the volume of 0.1 M titrant consumed was 7.8 ml, calculate the % purity of the sample.

- (A) 5.70%
- (B) 9.97%
- (C) 8.87%
- (D) 98.79%

Data for questions 74 and 75:

A drug, which is unstable to light, susceptible to oxygen and gets degraded in presence of metallic ions, has to be formulated in the form of a solution for injection.

74. Choose a suitable additive to improve the stability of the injection.

- (A) Preservative
- (B) Chelating agent
- (C) Buffer
- (D) Tonicity contributor

75. Select the appropriate filling and packing method for the above product.

- (A) Filling in an amber colored ampoule with an addition of antioxidant, replacing the inside air with nitrogen and sealing.
- (B) Filling with an antioxidant dissolved in the solution and sealing the ampoule.
- (C) Filling in an amber colored ampoule with a preservative and sealing.
- (D) Filling in an ampoule, sealing and giving directions to store it in dark.

Data for questions 76 and 77:

The usual adulterants for clove buds are clove stalks and thymol buds.

76. Clove stalks can be identified by the presence of

- (A) Starch grains
- (B) Cystoliths
- (C) Lignified sclereids
- (D) Acicular crystals of calcium oxalate

77. Anthrophylli can be identified by the presence of

- (A) Lignified sclereids
- (B) Acicular crystals of calcium oxalate
- (C) Cystoliths
- (D) Starch grains

Data for questions 78 to 80:

Plant tissue culture of carrot is being developed in the laboratory on a semisolid White's medium.

78. The micronutrient essential in the medium is:

- (A) NaCl
- (B) SnCl_2
- (C) KCl
- (D) CaCl_2

79. The pH of the medium is:

- (A) 6.6
- (B) 6.0
- (C) 5.6

(D) 5.0

80. The tissue growth observed is:

- (A) Undifferentiated cells suspended in the medium.
- (B) Undifferentiated cells in clusters distributed in the medium.
- (C) Differentiated mass of cells.
- (D) Surface growth of undifferentiated mass of cells.

Data for questions 81 and 82:

In glucose metabolism, name the enzymes catalyzing the following steps.

81. Conversion of glucose to glucose-6-phosphate.

- (A) Hexokinase
- (B) Glucose-6-phosphate dehydrogenase
- (C) Glycogen phosphorylase
- (D) Glycogen synthase

82. Conversion of 2-phosphoglycerate to phosphoenolpyruvate.

- (A) Pyruvate kinase
- (B) Phosphoglycerate mutase
- (C) Phosphoglycerate kinase
- (D) Enolase

Data for questions 83 and 84:

Methotrexate, Trimethoprim and Pyrimethamine are all known to be inhibitors of dihydrofolate reductase. Yet, they are classified in different therapeutic categories.

83. Trimethoprim has an advantage over methotrexate in its therapeutic category because

- (A) Trimethoprim binds to bacterial DHFR about 50,000 times more strongly as compared to the host DHFR.
- (B) Trimethoprim can be administered orally.
- (C) Trimethoprim exhibits no significant adverse effects.
- (D) Trimethoprim has additional anti-inflammatory properties.

84. Methotrexate is thought to exert its action by

- (A) Interfering with uric acid synthesis.
- (B) Intracellular formation of an amine adducts.
- (C) Forming a conjugate with nucleic acids.
- (D) Inhibiting the synthesis of folic acid.

Data for questions 85 to 87:

An administration officer having high blood pressure, gastric acidity and diabetes is prescribed enalapril, enalapril and tolbutamide.

85. From the structural features of the drugs, predict which will be ionized in the stomach.

- (A) Famotidine
- (B) Enalapril
- (C) Tolbutamide

(D) Enalapril and tolbutamide

86. The patient cannot tolerate enalapril. Which of the following can be substituted?

- (A) Omeprazole
- (B) Losartan
- (C) Rosiglitazone
- (D) Clofibrate

87. Famidine acts as

- (A) H1 — histamine antagonist
- (B) H2 — histamine antagonist
- (C) Proton pump inhibitor
- (D) H1 agonist

Data for questions 88 to 90:

2-Methoxy naphthalene on treatment with acetyl chloride in presence of AlCl₃ gives 2-acetyl-6-methoxy naphthalene. This is converted with a set of reagents X to 6-methoxy-2-naphthyl acetic acid, which is esterified with methanol to the methyl ester. Ester on treatment with Y gives DL-2-(6-methoxy-2-naphthyl)-propionic acid methyl ester. This on hydrolysis gives Z (final compound).

88. The set of reagents X are

- (A) Morpholine/Sulphur followed by H₂SO₄/H₂O
- (B) Morphine/Sulphur followed by HCl/H₂O
- (C) Formic acid/Cu followed by acetic acid
- (D) Hydroiodic acid followed by H₂SO₄/H₂O

89. Identify the reagents — Y.

- (A) NaOH/CH₃OH (B) NaH/CH₃I
- (C) Hydrazine/CH₃I (D) LiAlH₄/CH₃OH

90. The final compound Z is:

- (A) Naphazoline
- (B) Carprofen
- (C) Pranoprofen
- (D) Naproxen