

STATISTICS – III

Time Allowed : Three Hours

Maximum Marks : 200

**INSTRUCTIONS**

*Candidates should attempt FIVE questions in ALL including Questions No. 1 and 5 which are compulsory. The remaining THREE questions should be answered by choosing at least ONE question each from Section A and Section B.*

*The number of marks carried by each question is indicated against each.*

*Answers must be written only in ENGLISH.*

*(Symbols and abbreviations are as usual.)*

*If any data/value is to be assumed for answering a question, the same must be mentioned clearly.*

**SECTION A**

1. Answer any *five* parts : 8×5=40

- (a) Discuss Lahiri's method of selecting samples for PPS scheme. Give a suitable example to select 4 units when  $N = 10$ ,  $M = 60$ , by taking your choice of sample number and its corresponding size.

- (b) Explain the concept of simple random/sampling with and without replacement. In a simple random sampling, show that  $s^2$  is an unbiased estimator of  $S^2$ .
- (c) For the ratio estimator of the population mean  $\bar{Y}$ , obtain the bias of  $\bar{y}_r$  in terms of covariance. Further show that if the coefficient of variation of  $\bar{x}$  is sufficiently small, the bias compared to standard error of  $\bar{y}_r$  may be considered to be negligible.
- (d) For symmetric BIBD, show that  $|N| = \sqrt{r(r-\lambda)^{v-1}}$ , where  $N$  is the incidence matrix of SBIBD.
- (e) Discuss  $3^2$  factorial experiments. Explain a method to estimate all the main effects and interaction effects at single degree of freedom for  $3^2$  factorial experiment. Write its ANOVA table considering replication size 2.
- (f) Construct a key block of a  $3^4$  confounded factorial experiment into a block of size 9 by confounding the interactions  $ABC$  and  $AB^2D$ . Write all its generalized confounded interactions. Construct its one more block.

2. (a) Let for a stratified random sampling

$$n_i = \frac{W_i S_i}{\sqrt{\mu_0 c_i}}$$

where  $\mu_0$  is constant,  $c_i$  is the cost per unit in  $i^{\text{th}}$  stratum,  $W_i = \frac{N_i}{N}$ ,  $S_i^2$  is the mean square based on  $N_i$  units. Estimate the sample size  $n$  under optimum allocation for fixed cost  $c_0$ .

- (b) In two stage sampling with equal first stage units, obtain the variance of the sample mean  $\bar{\bar{y}}_2$ .
- (c) Discuss cluster sampling. Let  $\bar{\bar{y}}$  be the sample mean based on a sample of  $nM$  elements.  $\bar{y}$  is the sample mean based on a sample of  $nM$  elements drawn for SRS without replacement from  $NM$  elements in the population. Obtain Relative efficiency of  $\bar{\bar{y}}$  with respect to  $\bar{y}$  (without getting the derivation of  $V(\bar{y})$  and  $V(\bar{\bar{y}})$ ). Write its ANOVA table.
- (d) Explain aligned sample and unaligned systematic sampling method. Give an example of each with  $m = 3$ ,  $l = 3$ ,  $n = 3$  and  $k = 4$ , where  $nm$  units denotes systematic samples, provided units of population are arranged in the form of  $ml$  rows each containing  $nk$  units.  $10 \times 4 = 40$

3. (a) Discuss a method of construction of BIBD with parameters  $v = b = 11$ ,  $r = k = 6$  and  $\lambda = 3$ .
- (b) Explain Randomized block design. Obtain non-zero eigen value of C matrix of RBD. Estimate  $t_i$  ( $i = 1, 2, \dots, v$ ) using non-zero eigen value of C matrix.
- (c) Discuss missing plot techniques in a RBD. Suppose one observation, say,  $t_i$  is missing in one block of a RBD, derive a method to estimate that missing value.
- (d) Discuss symmetrical BIBD. For a SBIBD, show that any two blocks have exactly  $\lambda$  treatments in common. 10×4=40
4. (a) Explain difference estimator of population mean. Obtain its mean and variance. Hence obtain the linear regression estimator of  $\bar{Y}$ .
- (b) In a PPSWOR scheme, let  $z_i = \frac{y_i}{NP_i}$ . Obtain  $E(\bar{z})$  and  $V(\bar{z})$ .
- (c) Discuss group divisible PBIB design of two associate classes. Construct a PBIB design with parameters  $v = 8 = b$ ,  $r = 3 = k$ ,  $\lambda_1 = 0$  and  $\lambda_2 = 1$ . Further obtain the value of m and n.
- (d) Explain the layout of split plan designs. Write its model and assumptions. Give ANOVA table of sub-plot observations (only df and sum of squares). 10×4=40

## SECTION B

5. Attempt any *five* parts :

8×5=40

- (a) Explain moving average  $[m, p]$ . Let  $U_t = a_0 + a_1 t$ , with  $p = 1$  and  $m = 2k + 1$ , obtain the coefficient  $c_j$  of  $U_t$  and  $[m, p]$ .
- (b) Let  $U_t = a\xi + e_t$ ,  $-\infty < t < \infty$ , where  $e_t$ 's are i.i.d. with  $E(e_t) = 0$  and  $V(e_t) = 1$ . Show that the process is stationary with correlation.
- (c) For the following table :

**Commodity**

	A	B
$p_0$	1	1
$q_0$	10	5
$p_1$	2	X
$q_1$	5	2

where  $p$  and  $q$  stand for price and quantity for 0 and 1 time periods respectively. Find the value of  $X$  if the ratio between Laspeyres' (L) and Paasche's (P) index number is

$$L : P :: 28 : 27$$

- (d) For the general linear model  $Y = X\beta + e$ , following observations

$$\sum x_1^2 = 50, \quad \sum x_2^2 = 960, \quad \sum x_1 x_2 = -60,$$

$$\sum x_1 y = 30, \quad \sum x_2 y = 40, \quad \bar{X}_1 = 5, \quad \bar{X}_2 = 6 \quad \text{and} \\ \bar{Y} = 5$$

are given, where lower case letters  $y_j, x_j$  ( $j = 1, 2$ ) are given as  $y = Y - \bar{Y}$  and  $x_j = X_j - \bar{X}_j$ .

Obtain OLS estimator.

- (e) Discuss the effect of imperfect multicollinearity on tests and errors. Consider a model

$$y = \beta_0 + \beta_1 x_1 + \beta_2 x_2 + (e - \bar{e}) \text{ with}$$

$$\sum x_1 = \sum x_2 = 0 \text{ and } x_2 = \theta x_1 + v \text{ with}$$

$$\sum x_1^2 = \sum x_2^2 = 1, \quad \sum x_1 x_2 = \theta, \quad \sum v = 0 \text{ and}$$

$$\sum x_1 v = 0.$$

Do you think multicollinearity is present in the model? If yes, give reasons. Further show that as  $\theta$  increases,  $V(\hat{\beta}_1)$  also increases.

- (f) Explain demand function. Discuss Pigou's method of deriving demand curves from time series data. Indicate the assumptions made. Give a criticism of the method.

6. (a) Give the various steps for finding the variance of the random component using variate difference method. Is F-test used for testing the significance of homogeneity of two successive estimates of variance? If yes, ok, otherwise which test can be used? Discuss.
- (b) Explain the errors in the measurement of price and quantity index number.
- (c) Discuss second order autoregressive series. For this series, obtain complementary function (CF) only.
- (d) Explain curve of concentration. Obtain income concentration for Pareto's law of income distribution.

10×4=40

7. (a) For generalized least square linear model

$\underline{Y} = \underline{X}\underline{\beta} + \underline{e}$ , write assumptions. Obtain  $V(\hat{\underline{\beta}})$  and estimate  $\hat{\sigma}^2$  provided given another linear model is  $Y^* = X^* \beta + e^*$ , where  $X^* = T^{-1} X$ ,  $Y^* = T^{-1} Y$ ,  $e^* = T^{-1} e$ ,  $\Omega = TT'$  and  $e^* \sim N(0, \sigma^2)$ .

(b) If  $U = c x^\alpha y^\beta$  is an individual's utility function of two goods, show that the demand for the goods is

$$x = \frac{\alpha}{\alpha + \beta} \frac{\mu}{p_x} \quad \text{and} \quad y = \frac{\beta}{\alpha + \beta} \frac{\mu}{p_y}$$

where  $p_x$  and  $p_y$  are the fixed price and  $\mu$  be the individual fixed income.

(c) For which simultaneous equation, is indirect least square method of estimation used ? Discuss a method to estimate parameters using indirect least square estimator.

(d) Discuss economic forecasting of one single future observation on  $Y_T$ . 10×4=40

8. (a) For the auto-regressive scheme

$U_{t+2} + aU_{t+1} + bU_t = e_{t+2}$ ,  
show that if  $e$  is a random variable and the series is long, then

$$\frac{\text{Var}(U)}{\text{Var}(e)} = \frac{(1+b)}{(1-b)[(1+b)^2 - a^2]}$$

and hence show that, variance of the generated series may be much greater than that of  $e$  itself.

- (b) Explain the formulation of the problem of distribution of income. Further discuss its mathematical formulation in order to find number of persons with income  $x$  or more.
- (c) Discuss the method of estimating the parameters of simultaneous equation model using two stage least square method.
- (d) Explain rank condition of identification problem of simultaneous equation model.  $10 \times 4 = 40$